


Amendments to the Claims

This listing of the claims will replace all prior versions and listings of claims in the application:

Listing of Claims:

1. (Currently amended) A method of treatment for a mammal suffering from a dermatologic condition selected from the group consisting of contact dermatitis, acne, rosacea, and psoriasis, comprising administering a therapeutically effective amount of a compound of Formula I:



Formula I

wherein:

R¹ is: C(O)OR'; -C(O)NR'R"; -CH₂OR"'; cyano; optionally substituted heterocyclyl; optionally substituted heterocyclyl-alkyl; optionally substituted heteroaryl, or optionally substituted heteroaralkyl;

R² is: optionally substituted alkyl; optionally substituted cycloalkyl; optionally substituted aryl; optionally substituted aralkyl; optionally substituted heterocyclyl; optionally substituted heteroaryl; optionally substituted heteroaralkyl; an optionally substituted nucleoside; an optionally substituted amino acid; or an optionally substituted di-, tri- or tetra-peptide;

R³ is: optionally substituted alkyl; optionally substituted cycloalkyl; optionally substituted aryl; optionally substituted aralkyl; optionally substituted heterocyclyl; optionally substituted heteroaryl; optionally substituted heteroaralkyl; an optionally substituted nucleoside; an optionally substituted amino acid; or an optionally substituted di-, tri- or tetra-peptide;

R⁴ is: hydrogen; alkyl; alkylcarbonyl; (poly)alkoxyalkylene; or dialkoxyphosphoryloxy;

X is: —S—; —S(O)—; —S(O)₂—; or X taken together with R² is —P(O)(OR');

Y is: —S—; —S(O)—; —S(O)₂—; or Y taken together with R³ is —P(O)(OR')₂;

or X—R² taken together with Y—R³ form an optionally substituted aliphatic or aromatic ring;


R' is: hydrogen; alkenyl; optionally substituted alkyl; optionally substituted cycloalkyl; phosphoryl; or optionally substituted aryl;

R" is: hyrodgen; alkenyl; optionally substituted alkyl; or optionally substituted aryl;

or R' and R" together with the atom to which they are attached form a 5- to 7-membered aromatic, saturated or unsaturated ring, optionally incorporating one or more additional heteroatoms chosen from N, O, or S, and optionally substituted with one or more substitutents selected from the group consisting of optionally substituted lower alkyl, halo, cyano, alkylthio, lower alkoxy, carboxy, benzyl, and oxo;

R''' is: hydrogen; alkenyl; optionally substituted alkyl; acyl, optionally substituted cycloalkyl; phosphoryl; or optionally substituted aryl;

or a compound of Formula III:



Formula III

wherin:

R⁵ is: —C(O)OR^a; —C(O)NR^bR^b; —CH₂OR^d; —C(O)R^c; cyano; optionally substituted heterocycl; or optionally substituted heteroaryl;

R⁶ is: hydrogen; —C(O)OR^a; —C(O)NR^bR^b; —CH₂OR^d; —C(O)R^c; cyano; optionally substituted alkyl; optionally substituted heterocycl; optionally substituted aryl, or optionally substituted heteroaryl;

or R⁵ and R⁶ with the atom to which they are attached form an optionally substituted ring;

R⁷ is: optionally substituted alkyl; optionally substituted cycloalkyl; optionally substituted aryl; optionally substituted aralkyl; optionally substituted heterocycl; optionally substituted heteroaryl; optionally substituted heteroaralkyl; an optionally substituted nucleoside; an optionally substituted amino acid; or an optionally substituted di-, tri- or tetra-peptide; with the proviso that when R⁶ is alkyl, then R⁷ is optionally substituted heterocycl, optionally substituted heteroaryl, or optionally substituted heteroaralkyl,

or R⁵ and R⁷ with the atoms to which they are attached form an optionally substituted heterocyclic ring;

R⁸ is: hydrogen; alkyl; alkylcarbonyl; (poly)alkoxyalkylene; or dialkoxyphosphoryloxy;
Y' is: S—;

R^a is: hydrogen; alkenyl; optionally substituted alkyl; optionally substituted cycloalkyl;
or optionally substituted aryl;

R^b is: hydrogen; alkenyl; optionally substituted alkyl; or optionally substituted aryl;

or R^a and R^b together with the atom to which they are attached for a 5- to 7-membered aromatic, saturated or unsaturated ring, optionally incorporating one or more additional heteroatom chosen from N, O, or S, and optionally substituted with one or more substituents selected from the group consisting of optionally substituted lower alkyl, halo, cyano, alkylthio, lower alkoxy, carboxy, benzyl, and oxo;

R^c is: optionally substituted alkyl or optionally substituted aryl; and

R^d is: hydrogen; alkenyl; optionally substituted alkyl; acyl; optionally substituted cycloalkyl; or optionally substituted aryl; including single tautomers; single stereoisomers and mixtures of tautomers and/or stereoisomers, and the pharmaceutically acceptable salts thereof.

2. (Original) The method of claim 1, wherein X and Y are both —S—.

3. (Currently amended) A method of treatment for a mammal suffering from a dermatologic condition selected from the group consisting of contact dermatitis, acne, rosacea, and psoriasis comprising administering a therapeutically effective amount of a compound [[is]] selected from the group consisting of:

R-3-[2-(4-Amino-4-carboxy-butylamino)-2-(carboxymethyl-carbamoyl)-ethylsulfanyl]-2-[2-(4-amino-4-carboxy-butylamino)-2-(carboxymethyl-carbamoyl)-ethylsulfanylmethyl]-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(5-Chloro-benzothiazol-2-ylsulfanyl)-2-(5-chloro-benzothiazol-2-ylsulfanylmethyl)-4-hydroxy-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-Cyclohexylsulfanyl-2-cyclohexylsulfanylmethyl-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(pyrrolidine-1-carbothioylsulfanyl)-2-(pyrrolidine-1-carbothioylsulfanylmethyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(Benzoselenazol-2-ylsulfanyl)-2-(benzoselenazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(7-trifluoromethyl-quinolin-4-ylsulfanyl)-2-(7-trifluoromethyl-quinolin-4-ylsulfanylmethyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(2-sulfo-ethylsulfanyl)-2-(2-sulfo-ethylsulfanylmethyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

Hydroxy-5-oxo-3-(4-trifluoromethyl-pyrimidin-2-ylsulfanyl)-2-(4-trifluoromethyl-pyrimidin-2-ylsulfanylmethyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(1-oxy-pyridin-2-ylsulfanyl)-2-(1-oxy-pyridin-2-ylsulfanylmethyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(Benzothiazol-2-ylsulfanyl)-2-(benzothiazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2-Chloro-phenylsulfanyl)-2-(2-chloro-phenylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-Hexylsulfanyl-2-hexylsulfanylmethyl-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(1-phenyl-1*H*-tetrazol-5-ylsulfanyl)-2-(1-phenyl-1*H*-tetrazol-5-ylsulfanylmethyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(naphthalen-2-ylsulfanyl)-2-(naphthalen-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(4-phenyl-thiazol-2-ylsulfanyl)-2-(4-phenyl-thiazol-2-ylsulfanylmethyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-(5-sulfo-1*H*-benzoimidazol-2-ylsulfanyl)-5-(5-sulfo-1*H*-benzoimidazol-2-ylsulfanylmethyl)-3-hydroxy-5-hydroxymethyl-5*H*-furan-2-one;

3-(Furan-2-ylmethysulfanyl)-2-(furan-2-ylmethylsulfanylmethyl)-4-isobutyroloxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-[4-(2-methoxycarbonyl-vinyl)-phenylsulfanyl]-2-[4-(2-methoxycarbonyl-vinyl)-phenylsulfanylmethyl]-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(6-nitro-benzothiazol-2-ylsulfanyl)-2-(6-nitro-benzothiazol-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

2-(Benzothiazole-2-sulfinylmethyl)-3-(benzothiazol-2-ylsulfanyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2,4-Dichloro-benzylsulfanyl)-2-(2,4-dichloro-benzylsulfanyl)methyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(5-methoxy-benzothiazol-2-ylsulfanyl)-2-(5-methoxy-benzothiazol-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2-Chloro-6-fluoro-benzylsulfanyl)-2-(2-chloro-6-fluorobenzylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

Dimethylamino-acetic acid 3-(1*H*-benzoimidazol-2-ylsulfanyl)-2-(1*H*-benzoimidazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-ylmethyl ester;

3-(2-Chloro-4-fluoro-phenylsulfanyl)-2-(2-chloro-4-fluorophenylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(Benzothiazol-2-ylsulfanyl)-2-(benzothiazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid;

2-(Furan-2-ylmethanesulfinylmethyl)-3-(furan-2-ylmethanesulfonyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(furan-2-ylmethanesulfinyl)-2-(furan-2-ylmethanesulfinylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-(2,2-dimethyl-propionyloxy)-3-ethoxycarbonylmethylsulfanyl-2-ethoxycarbonylmethylsulfanyl)methyl-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2-Dimethylamino-ethylsulfanyl)-2-(2-dimethylamino-ethylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid;

3-(5,6-Dichloro-1*H*-benzoimidazol-2-ylsulfanyl)-2-(5,6-dichloro-1*H*-benzoimidazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-Benzylsulfanyl-2-benzylsulfanylmethyl-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-(1*H*-Benzoimidazol-2-ylsulfanyl)-3-hydroxy-1-oxa-7,9-diaza-spiro[4,5]dec-3-ene-2,6,8,10-tetraone;

3-(Benzothiazol-2-ylsulfanyl)-2-(benzothiazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid isopropyl ester;

4-Acetoxy-3-(furan-2-ylmethylsulfanyl)-2-(furan-2-ylmethylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Ethoxy-3-(1-ethyl-1*H*-benzoimidazol-2-ylsulfanyl)-2-(1-ethyl-1*H*-benzoimidazol-2-ylsulfanyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

2-(1*H*-Benzoimidazol-2-ylsulfanylmethyl)-4-ethoxy-3-(1-ethyl-1*H*-benzoimidazol-2-ylsulfanyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Acetoxy-3-benzylsulfanyl-2-benzylsulfanylmethyl-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(1-methyl-1*H*-benzoimidazol-2-ylsulfanyl)-2-(1-methyl-1*H*-benzoimidazol-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(1*H*-Benzothiazol-2-ylsulfanyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(5-methyl-1*H*-benzimidazol-2-ylsulfanyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid 2-isopropyl-5-methyl-cyclohexyl ester;

Di-[2-(4 Hydroxy-5-oxo-2-carboxylic acid methyl ester)]-disulfide; and

3-(4-Fluoro-benzylsulfanyl)-4-hydroxy-5-oxo-5*H*-furan-2,2-dicarboxylic acid diethyl ester.

4. (Currently amended) A method of treatment for a mammal suffering from a dermatologic condition selected from the group consisting of contact dermatitis, acne, rosacea, and psoriasis comprising administering a therapeutically effective amount of a compound [[is]] selected from the group consisting of:

3-(5-Chloro-benzothiazol-2-ylsulfanyl)-2-(5-chloro-benzothiazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-Cyclohexylsulfanyl-2-cyclohexylsulfanylmethyl-4-hydroxy-5-oxo-2,5,-dihydro-furan-2-carboxylic acid ethyl ester;

3-(Benzoselenazol-2-ylsulfanyl)-2-(benzoselenazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-fur-an-2-carboxylic acid ethyl ester;

3-(Benzothiazol-2-ylsulfanyl)-2-(benzothiazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2-Chloro-phenylsulfanyl)-2-(2-chloro-phenylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-Hexylsulfanyl-2-hexylsulfanylmethyl-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(naphthalen-2-ylsulfanyl)-2-(naphthalen-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(4-phenyl-thiazol-2-ylsulfanyl)-2-(4-phenyl-thiazol-2-ylsulfanylmethyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(Furan-2-ylmethylsulfanyl)-2-(furan-2-ylmethylsulfanylmethyl)-4-isobutyryloxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2,4-Dichloro-benzylsulfanyl)-2-(2,4-dichloro-benzylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2-chloro-4-fluoro-phenylsulfanyl)-2-(2-chloro-4-fluoro-phenylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(furan-2-ylmethanesulfinyl)-2-(furan-2-ylmethanesulfinylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-(2,2-dimethyl-propionyloxy)-3-ethoxycarbonylmethylsulfanyl-2-ethoxycarbonylmethylsulfanylmethyl-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(Benzothiazol-2-ylsulfanyl)-2-(benzothiazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid isopropyl ester;

4-Hydroxy-3-(1-methyl-1*H*-benzoimidazol-2-ylsulfanyl)-2-(1-methyl-1*H*-benzoimidazol-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(1*H*-Benzoimidazol-2-ylsulfanyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3(5-methyl-1*H*-benzoimidazol-2-ylsulfanyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid 2-isopropyl-5-methyl-cyclohexyl ester;

Di-[2-(4-Hydroxy-5-oxo-2-carboxylic acid methyl ester)]-disulfide; and

3-(4-Fluoro-benzylsulfanyl)-4-hydroxy-5-oxo-5*H*-furan-2,2,-dicarboxylic acid diethyl ester.

5. (Canceled)

6. (Original) The method of claim 1, comprising topically administering at least one compound of Formula I or of Formula III.

7. (Currently amended) The method of claim 3, comprising applying to the area of skin in need of such treatment a composition comprising at least one compound selected from the group consisting of:

R-3-[2-(4-Amino-4-carboxy-butyrylamino)-2-(carboxymethyl-carbamoyl)-ethylsulfanyl]-2-[2-(4-amino-4-carboxy-butyrylamino)-2-(carboxymethyl-carbamoyl)-ethylsulfanylmethyl]-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(5-Chloro-benzothiazol-2-ylsulfanyl)-2-(5-chloro-benzothiazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-Cyclohexylsulfanyl-2-cyclohexylsulfanylmethyl-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(pyrrolidine-1-carbothioylsulfanyl)-2-(pyrrolidine-1-carbothioylsulfanyl methyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(Benzoselenazol-2-ylsulfanyl)-2-(benzoselenazol-2-ylsulfanyl methyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(7-trifluoromethyl-quinolin-4-ylsulfanyl)-2-(7-trifluoromethyl-quinolin-4-ylsulfanyl methyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(2-sulfo-ethylsulfanyl)-2-(2-sulfo-ethylsulfanyl methyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(4-trifluoromethyl-pyrimidin-2-ylsulfanyl)-2-(4-trifluoromethyl-pyrimidin-2-ylsulfanyl methyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(1-oxy-pyridin-2-ylsulfanyl)-2-(1-oxy-pyridin-2-ylsulfanyl methyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(Benzothiazol-2-ylsulfanyl)-2-(benzothiazol-2-ylsulfanyl methyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2-Chloro-phenylsulfanyl)-2-(2-chloro-phenylsulfanyl methyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-Hexylsulfanyl-2-hexylsulfanyl methyl-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(1-phenyl-1*H*-tetrazol-5-ylsulfanyl)-2-(1-phenyl-1*H*-tetrazol-5-ylsulfanyl methyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(naphthalen-2-ylsulfanyl)-2-(naphthalen-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-5-oxo-3-(4-phenyl-thiazol-2-ylsulfanyl)-2-(4-phenyl-thiazol-2-ylsulfanylmethyl)-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-(5-sulfo-1*H*-benzoimidazol-2-ylsulfanyl)-5-(5-sulfo-1*H*-benzoimidazol-2-ylsulfanylmethyl)-3-hydroxy-5-hydroxymethyl-5*H*-furan-2-one;

3-(Furan-2-ylmethylsulfanyl)-2-(furan-2-ylmethylsulfanylmethyl)-4-isobutyryloxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-[4-(2-methoxycarbonyl-vinyl)-phenylsulfanyl]-2-[4-(2-methoxycarbonyl-vinyl)-phenylsulfanylmethyl]-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(6-nitro-benzothiazol-2-ylsulfanyl)-2-(6-nitro-benzothiazol-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

2-(Benzothiazole-2-fulfinylmethyl)-3-(benzothiazol-2-ylsulfanyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2,4-Dichloro-benzylsulfanyl)-2-(2,4,-dichloro-benzylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(5-methoxy-benzothiazol-2-ylsulfanyl)-2-(5-methoxy-benzothiazol-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2-Chloro-6-fluoro-benzylsulfanyl)-2-(2-chloro-6-fluoro-benzylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

Dimethylamino-acetic acid 3-(1*H*-benzimidazol-2-ylsulfanyl)-2-(1*H*-benzimidazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-ylmethyl ester;

3-(2-Chloro-4-fluoro-phenylsulfanyl)-2-(chloro-4-fluoro-phenylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(Benzothiazol-2-ylsulfanyl)-2-(benzothiazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid;

2-(Furan-2-ylmethanesulfinylmethyl)-3-(furan-2-ylmethanesulfonyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(furan-2-ylmethanesulfinyl)-2-(furan-2-ylmethanesulfinylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-(2,2-dimethyl-propionyloxy)-3-ethoxycarbonylmethylsulfanyl-2-ethoxycarbonylmethylsulfanylmethyl-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(2-Dimethylamino-ethylsulfanyl)-2-(2-dimethylamino-ethylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid;

3-(5,6-Dichloro-1*H*-benzimidazol-2-ylsulfanyl)-2-(5,6-dichloro-1*H*-benzimidazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-Benzylsulfanyl-2-benzylsulfanylmethyl-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-(1*H*-Benzoimidazol-2-ylsulfanyl)-3-hydroxy-1-oxa-7,9-diaza-spiro[4,5]dec-3-ene-2,6,8,10-tetraone;

3-(Benzothiazol-2-ylsulfanyl)-2-(benzothiazol-2-ylsulfanylmethyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid isopropyl ester;

4-Acetoxy-3-(furan-2-ylmethylsulfanyl)-2-(furan-2-ylmethylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Ethoxy-3-(1-ethyl-1*H*-benzoimidazol-2-ylsulfanyl)-2-(1-ethyl-1*H*-benzoimidazol-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

2-(1*H*-Benzoimidazol-2-ylsulfanylmethyl)-4-ethoxy-3-1-ethyl-1*H*-benzoimidazol-2-ylsulfanyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Acetoxy-3-benzylsulfanyl-2-benzylsulfanylmethyl-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(1-methyl-1*H*-benzoimidazol-2-ylsulfanyl)-2-(1-methyl-1*H*-benzoimidazol-2-ylsulfanylmethyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

3-(1*H*-Benzoimidazol-2-ylsulfanyl)-4-hydroxy-5-oxo-2,5-dihydro-furan-2-carboxylic acid ethyl ester;

4-Hydroxy-3-(5-methyl-1*H*-benzoimidazol-2-ylsulfanyl)-5-oxo-2,5-dihydro-furan-2-carboxylic acid 2-isopropyl-5-methyl-cyclohexyl ester;

Di-[2-(4-Hydroxy-5-oxo-2-carboxylic acid methyl ester)] disulfide; and

3-(4-Fluoro-benzylsulfanyl)-4-hydroxy-5-oxo-5H-furan-2,2-dicarboxylic acid diethyl ester.

8. (Previously Presented) The method of claim 1, comprising administering a formulation for topical application comprising one or more topical excipients and at least one compound of Formula I or of Formula III.

9. (Currently amended) The method of claim 1, additionally comprising an additional benefit agent, selected from the group consisting of sunscreens, retinoid and derivatives thereof, antioxidants, hydroxyacids, botanical extracts, salicylic acid, benzoyl peroxide, antibiotics, antiandrogens, anti-inflammatory agents, vitamins, tocopherol (α -, β -, γ -, δ -) and esters thereof, corticosteroids ~~corticosteroid~~ and mixtures thereof.

10. (Currently amended) The method of claim 3, additionally comprising an additional benefit agent, selected from the group consisting of sunscreens, retinoid and derivatives thereof, antioxidants, hydroxyacids, botanical extracts, salicylic acid, benzoyl peroxide, antibiotics, antiandrogens, anti-inflammatory agents, vitamins, tocopherol (α -, β -, γ -, δ -) and esters thereof, corticosteroids ~~corticosteroid~~ and mixtures thereof.

11. (Previously Presented) The method of claim 3, comprising administering a formulation for topical application comprising one or more topical excipients and at least one compound of Formula I or of Formula III.

12. (Canceled)

13. (Currently amended) The method of claim 4, additionally comprising an additional benefit agent, selected from the group consisting of sunscreens, retinoid and derivatives thereof, antioxidants, hydroxyacids, botanical extracts, salicylic acid, benzoyl peroxide, antibiotics, antiandrogens, anti-inflammatory agents, vitamins, tocopherol (α -, β -, γ -, δ -) and esters thereof, corticosteroids ~~corticosteroid~~ and mixtures thereof.

14. (Previously Presented) The method of claim 4, comprising administering a formulation for topical application comprising one or more topical excipients and at least one compound of Formula I or of Formula III.